

Two Theorems by Helmholtz

H.Fleming

Instituto de Física, Universidade de São Paulo

November 17, 2001

Abstract

Why is it that *div* and *curl* crop up everywhere in the study of vector fields? These theorems by Helmholtz explain.

Por que os operadores *div* e *rot* aparecem em toda parte, no estudo de campos vetoriais? Estes teoremas de Helmholtz explicam.

1 Introduction

What is usually called Helmholtz theorem[1] is the fact that a vector field \vec{V} which vanishes at infinity is completely determined by giving, everywhere, $div\vec{V}$ and $curl\vec{V}$. Helmholtz arrived at it in his study of vortices in fluids[4]. Nowadays it is mostly used in Electrodynamics, where the very Maxwell equations express its importance by being nothing else than the specification of what the *div* and *curl* of the electric and magnetic fields are.

2 The First Theorem

Actually, the Helmholtz theorem¹ proves a slightly different thing, from which the statement above follows immediately: a vector field \vec{V} which vanishes at the boundaries can be written as the sum of two terms, one of which is

¹In modern form, much generalized, this is the famous Kodaira-Hodge-De Rham decomposition theorem[6]

irrotational and the other, solenoidal (that is, divergenceless)[2]. Consider the following well-known identity for an arbitrary vector field $\vec{Z}(\vec{r})$:

$$-\vec{\nabla}^2 \vec{Z} = -\vec{\nabla}(\vec{\nabla} \cdot \vec{Z}) + \vec{\nabla} \times \vec{\nabla} \times \vec{Z} \quad (1)$$

If we now take our vector field to be

$$\vec{V} = -\vec{\nabla}^2 \vec{Z} \quad (2)$$

then it follows that

$$\vec{V} = -\vec{\nabla}U + \vec{\nabla} \times \vec{W} \quad (3)$$

with

$$U = \vec{\nabla} \cdot \vec{Z} \quad (4)$$

and

$$\vec{W} = \vec{\nabla} \times \vec{Z} \quad (5)$$

Eq.(3) is Helmholtz's theorem, as $\vec{\nabla}U$ is irrotational and $\vec{\nabla} \times \vec{W}$ is solenoidal.

But, is it general? It assumes that our vector field can be written as the Laplacian of some other one. . .

This constitutes, however, no problem if \vec{V} vanishes at infinity fast enough, for, then, the equation

$$\vec{\nabla}^2 \vec{Z} = -\vec{V} , \quad (6)$$

which is Poisson's equation, has always the solution

$$\vec{Z}(\vec{r}) = \frac{1}{4\pi} \int d^3 r' \frac{\vec{V}(\vec{r}')}{|\vec{r} - \vec{r}'|} . \quad (7)$$

It is now a simple matter to prove, from Eq.(3), that \vec{V} is determined from its *div* and *curl*. Taking, in fact, the divergence of Eq.(3), we have:

$$\text{div} \vec{V} = -\vec{\nabla}^2 U \quad (8)$$

which is, again, Poisson's equation, and, so, determines U as

$$U(\vec{r}) = \frac{1}{4\pi} \int d^3 r' \frac{\vec{\nabla}' \cdot \vec{V}(\vec{r}')}{|\vec{r} - \vec{r}'|} \quad (9)$$

Take now the *curl* of Eq.(3). We have

$$\begin{aligned} \vec{\nabla} \times \vec{V} &= \vec{\nabla} \times \vec{\nabla} \times \vec{W} \\ &= \vec{\nabla}(\vec{\nabla} \cdot \vec{W}) - \vec{\nabla}^2 \vec{W} \end{aligned} \quad (10)$$

Now, $\vec{\nabla} \cdot \vec{W} = 0$, as $\vec{W} = \vec{\nabla} \times \vec{Z}$, so another Poisson equation determines \vec{W} . Using U and \vec{W} so determined in Eq.(3) proves our contention.

3 Applications

We start with a very simple one. Consider a homogeneous, ohmic conductor, and let the current density \vec{j} vanish at the closed surface which encircles it. We want to inquire as to the possibility that a stationary current may exist in this conductor. We then have $\vec{\nabla} \cdot \vec{j} = 0$, as the current is supposed to be stationary, and $\vec{j} = \sigma \vec{E}$, which is Ohm's law. Also, $\vec{\nabla} \times \vec{E} = 0$, for a static \vec{E} . But then,

$$\vec{\nabla} \times \vec{j} = \sigma \vec{\nabla} \times \vec{E} = 0 \quad (11)$$

and so we have both $\vec{\nabla} \cdot \vec{j} = 0$ and $\vec{\nabla} \times \vec{j} = 0$. Besides, \vec{j} vanishes at the boundaries. It follows, then, from Helmholtz's theorem, that $\vec{j} = 0$. So, no stationary current can run under these conditions. Notice that this is true also for a torus and its continuous deformations, so that it applies to any closed circuit, proving the necessity that the condition $\vec{j} = \sigma \vec{E}$ be broken somewhere in the circuit (where the electromotive force is located).

Consider now the electromagnetic potentials. The Maxwell equation

$$\vec{\nabla} \cdot \vec{B} = 0 \quad (12)$$

states that there exists a vector field \vec{A} such that

$$\vec{B} = \vec{\nabla} \times \vec{A} \quad (13)$$

this \vec{A} being called the vector potential. We may assume that \vec{A} vanishes at infinity. Now, what we know about \vec{A} is just its *curl*. Therefore, by Helmholtz's theorem, we are free to choose the value of its divergence. For instance, we may take $\vec{\nabla} \cdot \vec{A} = 0$, determining completely \vec{A} . This is the so-called Coulomb gauge. But we can also put $\vec{\nabla} \cdot \vec{A} = -\frac{1}{c} \frac{\partial \phi}{\partial t}$, ϕ being the scalar potential. This is the Lorentz gauge.

4 The Second Theorem

There is another important result by Helmholtz in the same paper we mentioned above. It concerns the most general infinitesimal deformation of a plastic (that is, non-rigid) body.²

²A beautiful treatment of this topic, as well as of the whole vector analysis, is found in Sommerfeld's *Mechanics of Deformable Bodies*.^[5]

Consider a small volume element of a deformable body (a fluid, say). Putting the origin of the coordinates inside this volume element, let \vec{r} be the position of a generic point P . The origin is denoted by O . After a deformation, the material point P has a new position vector, $\vec{r} + \vec{s}$. Also the point of the body located at the origin moved, its position now being given by \vec{s}_0 . We assume, as physically reasonable, that \vec{s} is, as a function of position, continuous and differentiable to any order. In what follows, in order to get results of enough generality, we assume P very close to O , that is to say, we consider an infinitesimal volume of the body. Notice that \vec{s} itself doesn't have to be small in what follows, though, in considering deformations, this is somewhat academic, as finite deformations can always be obtained from infinitesimal ones (for instance, by Lie group methods). This notwithstanding, the geometrical interpretation we will get is only clear for infinitesimal \vec{s} . On the other hand, the application for the electric field which is done below would be impoverished by confining \vec{s} to be infinitesimal.

A Taylor expansion for \vec{s} gives

$$\vec{s}(\vec{r}) = \vec{s}_0 + (\vec{r} \cdot \vec{\nabla}) \vec{s} \quad (14)$$

neglecting further corrections, what is allowed by the restriction to infinitesimal volume. Here we wrote \vec{s}_0 for $\vec{s}(\vec{0})$. In more detail, if s_i is the i -th component of \vec{s} , one has

$$s_i(\vec{r}) = (s_0)_i + \sum_l x_l \left(\frac{\partial s_i}{\partial x_l} \right)_{\vec{r}=0} \quad (15)$$

This can be abbreviated to

$$s_i = (s_0)_i + x_l \partial_l s_i \quad (16)$$

or,

$$s_i = (s_0)_i + x_l T_{li} \quad (17)$$

where, obviously,

$$T_{li} = (\partial_l s_i)_{\vec{r}=0} . \quad (18)$$

In order to analyse this deformation in terms of more basic ones, let us decompose T_{li} in the following way:

$$T_{li} = \frac{1}{2}(T_{li} + T_{il}) + \frac{1}{2}(T_{li} - T_{il}) \quad (19)$$

and consider separately the symmetric and the antisymmetric parts. The antisymmetric part is

$$\frac{1}{2}(\partial_l s_i - \partial_i s_l) = \frac{1}{2}\epsilon_{lik}(\text{curl } \vec{s})_k . \quad (20)$$

The symmetric part is $S_{li} \equiv \frac{1}{2}(T_{li} + T_{il})$. It can be decomposed as follows:

$$S_{li} = \frac{1}{3}\delta_{li}S + (S_{li} - \frac{1}{3}\delta_{li}S) \quad (21)$$

where $S \equiv \text{Tr}(S_{li}) = S_{ii}$. Now, $S_{li} = \frac{1}{2}(\partial_l s_i + \partial_i s_l)$, so that

$$S = \partial_i s_i = \vec{\nabla} \cdot \vec{s} . \quad (22)$$

We can therefore write Eq.(21) as

$$S_{li} = \frac{1}{3}\delta_{li}\vec{\nabla} \cdot \vec{s} + S_{li}^0 \quad (23)$$

where S_{li}^0 is a traceless symmetric tensor. Going now to Eq.(17), we can write

$$s_i = (s_0)_i + x_l \frac{1}{2}\epsilon_{lik}(\text{curl } \vec{s})_k + \frac{1}{3}x_i \text{div } \vec{s} + x_l S_{li}^0 \quad (24)$$

or,

$$\vec{s} = \vec{s}_0 + \frac{1}{2}(\text{curl } \vec{s} \times \vec{r}) + \frac{1}{3}\vec{r}\text{div } \vec{s} + \vec{r} \cdot \mathbf{S}^0 \quad (25)$$

where the last term is a vector whose i-th component is $x_l S_{li}^0$. An object like \mathbf{S}^0 is sometimes called a *dyadic*. It is a second-order tensor. Let us examine the meaning of the several terms of Eq.(25). The first term of the second member is a translation. The second is an infinitesimal rotation around the axis $(\text{curl } \vec{s})_{\vec{r}=0}$, and the remaining terms describe a dilatation of the volume element. To understand them better, let us suppose temporarily that the translation and the rotation vanish, so that we have, for a component of \vec{s} :

$$s_i = \frac{1}{3}\text{div } \vec{s} x_i + x_l S_{li}^0 . \quad (26)$$

Now, S_{li}^0 is a symmetric matrix, so it can be diagonalized. This means that we can change coordinate axes in such a way that, in the new ones, the

matrix elements of S^0 have the form $S_{li}^0 = \delta_{li}S_{ii}^0$, where no sum is implied in this last expression. So, if the new coordinates are denoted by X_i , we have

$$\sum_l x_l S_{li}^0 = \sum_l X_l \delta_{li} S_{ii}^0 = X_i S_{ii}^0 \quad \text{no sum on } i \quad (27)$$

so that

$$s_i = \frac{1}{3} \text{div } \vec{s} X_i + S_{ii}^0 X_i \quad (\text{no sum}) \quad (28)$$

Consider the infinitesimal ‘‘cube’’ centered at O , with P as a vertex. (The ‘‘cube’’ may have curvilinear edges, at least after the deformation). Its volume before the deformation was, in the new coordinates, $V = X_1 X_2 X_3$. After the deformation, it is $V' = (X_1 + s_1)(X_2 + s_2)(X_3 + s_3)$. As we have, considering only the dilatation,

$$s_i = \frac{1}{3} \text{div } \vec{s} X_i + S_{ii}^0 X_i = X_i \left(\frac{1}{3} \text{div } \vec{s} + S_{ii}^0 \right) \quad (29)$$

then

$$X_i + s_i = X_i \left(1 + \frac{1}{3} \text{div } \vec{s} + S_{ii}^0 \right) \equiv X_i (1 + \epsilon_i) \quad (30)$$

and, for the volume,

$$V' = X_1 X_2 X_3 (1 + \epsilon_1)(1 + \epsilon_2)(1 + \epsilon_3) \quad (31)$$

which, to first order is

$$V' = V(1 + \epsilon_1 + \epsilon_2 + \epsilon_3) \quad (32)$$

that is,

$$V' = V(1 + \text{div } \vec{s} + S_{11}^0 + S_{22}^0 + S_{33}^0) \quad (33)$$

but, as S_{ij}^0 is traceless,

$$V' = V(1 + \text{div } \vec{s}) . \quad (34)$$

So, the S_{ij}^0 do not contribute to the change of volume, but do contribute to the change of the form of the little cube. Referring again to Eq.(25), we see that \vec{s} is written as a sum of a translation, plus a rotation, plus an isotropic dilatation, plus a volume-conserving deformation.³

³This is, in fact, the decomposition, following Weyl[3], of the tensor T_{li} into its irreducible components under rotations. This corresponds to analysing an infinitesimal deformation in terms of quantities which transform as irreducible representations of the rotation group. In particular, the traceless symmetric tensor ‘‘carries’’ an irreducible representation of dimension 5, an anisotropic object.

Notice that Eq.(25) was obtained using only Calculus. So, it should apply to any vector field whatsoever. To shed more light on the role of each term of the expansion, let us use it for the electric field. We then have:

$$\vec{E}(\vec{r}) = \vec{E}(\vec{0}) + \frac{\vec{r}}{3}(\text{div } \vec{E})_0 + \frac{1}{2}[(\text{curl } \vec{E})_0 \times \vec{r}] + \vec{r} \cdot \mathbf{S}^0 . \quad (35)$$

As $(\text{div } \vec{E})_0 = 4\pi\rho(\vec{0})$, the second term reads

$$\frac{\vec{r}}{3}(\text{div } \vec{E})_0 = \frac{\vec{r}}{3}4\pi\rho(\vec{0}) = \frac{4\pi}{3}r^3\rho(\vec{0})\frac{\vec{r}}{r^3} . \quad (36)$$

Now, this is the field of a uniformly charged sphere of radius r at a point of its surface. It is a radial field to be added to \vec{E}_0 .

The second term vanishes if the fields are static. Otherwise,

$$(\text{curl } \vec{E})_0 = -\frac{1}{c}\left(\frac{\partial \vec{B}}{\partial t}\right)_0 \quad (37)$$

and

$$\frac{1}{2}[(\text{curl } \vec{E})_0 \times \vec{r}] = -\frac{1}{2c}\frac{\partial}{\partial t}[\vec{B}(\vec{0}) \times \vec{r}] . \quad (38)$$

Consider a uniform magnetic field of value $\vec{B}(\vec{0})$. A vector potential corresponding to it is

$$\vec{A}(\vec{r}) = \frac{1}{2}\vec{B}(\vec{0}) \times \vec{r} \quad (39)$$

so that we have

$$\frac{1}{2}[(\text{curl } \vec{E})_0 \times \vec{r}] = -\frac{1}{c}\frac{\partial \vec{A}}{\partial t} . \quad (40)$$

This means that the third term of (35) is the contribution of the magnetic field at the origin, treated as follows: extend the value of \vec{B} at $\vec{0}$ to a uniform field. Compute its vector potential and then add the term $-\frac{1}{c}\frac{\partial \vec{A}}{\partial t}$ to $\vec{E}(\vec{0})$. All the rest of the field at \vec{r} (and that could be a lot!) comes from the term \mathbf{S}^0 .

5 Conclusion

The first Helmholtz theorem explains why the *div* and *curl* are so ubiquitous in vector fields. It also explains the structure of Maxwell equations, usually

written in such a way that the first member of each is either a *div* or a *curl*. The second explains the meanings, and so also the names, of those operators. Considering that Hermann von Helmholtz graduated in Medicine, not in Physics, that was not a bad job at all!

I wish to thank Walter F. Wreszinski for enlightening comments.

References

- [1] W.K.H. Panofsky and M. Phillips, *Classical Electricity and Magnetism*, New York, 1970.
- [2] D.H. Kobe, Am. J. Phys. **54**, 552(1985).
- [3] H. Weyl, *The Classical Groups*, Princeton.
- [4] H.v. Helmholtz, *Über Integrale der hydrodynamischen Gleichungen, welche den Wirbelbewegungen entsprechen*, Crelles J. **55**,25(1858).
- [5] A. Sommerfeld, *Mechanics of Deformable Bodies* New York (1959).
- [6] R. Aldrovandi, J.G. Pereira, *An Introduction to Geometrical Physics*, World Scientific (1995).